



PREMIUM FEED SUPPLEMENTS

1505 4th Ave East, Horton, KS 66439
ph 785-486-2626 Fax 785-486-2244

Range MaXX

Protein	25.00% min	Cobalt	4 ppm
NPN	8.00% min	Copper	600 ppm
Crude Fat	3.50% min	Iodine	8 ppm
Crude Fiber	5.00% min	Manganese	750 ppm
Calcium	4.75% min 5.25% max	Selenium	5 ppm
Phosphorus	1.00% min	Zinc	1,200 ppm
Salt	13.00% min 14.00% max	Vitamin A	100,000 IU/lb
Magnesium	4.00% min	Vitamin D	20,000 IU/lb
Potassium	1.00% min	Vitamin E	80 IU/lb
Sulfur	2.25% min		

Ingredient Deck:

Corn Distillers Dried Grains with Solubles, Corn Condensed Distillers Solubles, Salt, Calcium Carbonate, Ammonium Sulfate, Magnesium Oxide, Mono-Calcium Phosphate, Corn Gluten Meal, Grain Product, Biuret, Calcium Sulfate, Calcium Hydroxide, Potassium Chloride, Yeast Culture, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Zinc Polysaccharide Complex, Zinc Protein, Manganese Polysaccharide Complex, Manganese Protein, Copper Polysaccharide Complex, Copper Protein, Aspergillus Oryzae Extract, Aspergillus Niger Fermentation Liquid, Cobalt Carbonate, Sodium Selenite, Ethylenediamine Dihydroiodide, Vitamin A Acetate (Stability Improved), Vitamin D3 Supplement, Vitamin E Supplement.

Feeding Directions:

1. Before feeding SweetPro® Range MaXX lick blocks for the first time, animals should be well fed.
2. Provide access for animals to have plenty of clean, fresh water at all times.
3. Provide an adequate source of good quality roughage and forage at all times (all they can consume).
4. Have one lick block available per 25 head of beef cattle.
5. Place lick blocks near water sources or areas cattle frequent regularly.
6. Make lick blocks available in more than one location in large pastures or where more than one water source is available.
7. Other salt and minerals may be provided free choice to balance specific needs.
8. A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended range of intake is .50 to .75 lbs. per head per day.